



[PubMed](#)
[Nucleotide](#)
[Protein](#)
[Genome](#)
[Structure](#)
[PopSet](#)
[Taxonomy](#)
[OMIM](#)

Search  for

[Limits](#)
[Preview/Index](#)
[History](#)
[Clipboard](#)
[Details](#)

Entrez

Entrez Version

Entrez PubMed

Overview

Help/FAQ

Feedback

New/Noteworthy

PubMed Services

Journal Browser

Med E Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cuba

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

Clinical Trials.gov

PubMed Central

Privacy Policy

☐ 1: EMBO J 1996 May 15;15  
 (10):2371-80

[Related Articles](#)
[Nucleotide](#)
[Protein](#)
[Books](#)
[LinkOut](#)

## WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na<sup>+</sup> channel deleted in Liddle's syndrome.

**Staub O, Dho S, Henry P, Correa J, Ishikawa T, McGlade J, Rotin D.**

The Hospital For Sick Children, Division of Respiratory Research, Toronto, Ontario, Canada.

The amiloride-sensitive epithelial sodium channel (ENaC) plays a major role in sodium transport in kidney and other epithelia, and in regulating blood pressure. The channel is composed of three subunits (alpha, beta, gamma) each containing two proline-rich sequences (P1 and P2) at its C-terminus. The P2 regions in human beta and gammaENaC, identical to the rat betagammaENaC, were recently shown to be deleted in patients with Liddle's syndrome (a hereditary form of hypertension), leading to hyperactivation of the channel. Using a yeast two-hybrid screen, we have now identified the rat homologue of Nedd4 (rNedd4) as the binding partner for the P2 regions of beta and gammaENaC. rNedd4 contains a Ca<sup>2+</sup> lipid binding (CaLB or C2) domain, three WW domains and a ubiquitin ligase (Hect) domain. Our yeast two-hybrid and in vitro binding studies revealed that the rNedd4-WW domains mediate this association by binding to the P2 regions, which include the PY motifs (XPPXY) of either betaENaC (PPPNY) or gammaENaC (PPPRY). SH3 domains were unable to bind these sequences. Moreover, mutations to Ala of Pro616 or Tyr618 within the betaENaC P2 sequence (to PPANY or PPPNA, respectively), recently described in Liddle's patients, led to abrogation of rNedd4-WW binding. Nedd4-WW domains also bound to the proline-rich C-terminus (containing the sequence PPPAY) of alphaENaC, and endogenous Nedd4 co-immunoprecipitated with alphaENaC expressed in MDCK cells. These results demonstrate that the WW domains of rNedd4 bind to the PY motifs deleted from beta or gammaENaC in Liddle's syndrome patients, and suggest that Nedd4 may be a regulator (suppressor) of the epithelial Na<sup>+</sup> channel.

PMID: 8665844 [PubMed - indexed for MEDLINE]